arXiv:1704.03637v1 [math.NT] 12 Apr 2017

Degrees of irreducible polynomials over binary field

Yaotsu Chang * Chong-Dao Lee† Chia-an Liu‡

April 13, 2017

Abstract

An algorithm for factoring polynomials over finite fields is given by Berlekamp in 1967. The main tool was the matrix Q corresponding to each polynomial. This paper studies the degrees of polynomials over binary field that associated with their corresponding matrices Q and irreducibility.

Keywords: Irreducible polynomial, binary field, Berlekamp matrix.

MSC2010: 11T06.

1 Introduction

Let $F = \mathbb{F}_2 = \{0, 1\}$ be the binary finite field. Then for each polynomial $f(x) \in F[x]$ of degree m, the *Berlekamp matrix* Q proposed in [1] of $f(x)$ is the $m \times m$ matrix over F whose ith row represents $x^{2(i-1)}$ reduced modulo $f(x)$. Specifically,

$$x^{2i} \equiv \sum_{j=0}^{m-1} Q_{i+1,j+1} x^j \pmod{f}$$

for $i = 0, 1, \ldots, m-1$.

*Department of Financial and Computational Mathematics, I-Shou University, Taiwan R.O.C..

†Department of Communication Engineering, I-Shou University, Taiwan R.O.C..

‡Corresponding author. E-mail address: liuciaan8@gmail.com, Department of Financial and Computational Mathematics, I-Shou University, Taiwan R.O.C..

2 Preliminary

Let G be the Berlekamp matrix with respect to the polynomial $f(x)$ of degree m over F. It is not difficult to show that $G^m = I_m$ if and only if $f(x)$ has no square factors, where I_m denotes the identity matrix of order m. If $f(x)$ has no square factors then the *order* $o(f(x)) = o(G)$ of $f(x)$ is defined to be the least positive integer such that

$$G^{o(G)} = I_m.$$

Lemma 2.1. *Let $f(x) = \prod_{i=1}^{r} g_i(x)$ where $g_i(x)$ are distinct polynomials of order d_i over F for $1 \leq i \leq r$. Then*

$$o(G) = lcm(d_1, d_2, \ldots, d_r).$$

It is not hard to have the following observation for least common multiple.

Lemma 2.2. *For positive integers $n_1, n_2, \ldots, n_\ell$ and k, the least common multiple*

$$lcm(kn_1, kn_2, \ldots, kn_\ell) = k \cdot lcm(n_1, n_2, \ldots, n_\ell).$$

Proof. Let $A = \mathrm{lcm}(kn_1, kn_2, \ldots, kn_\ell)$ and $B = \mathrm{lcm}(n_1, n_2, \ldots, n_\ell)$. For each $i = 1, 2, \ldots, \ell$, one has $kn_i | kB$ since $n_i | B$, and hence $A | kB$. On the other hand, $kn_i | A$ implies $n_i | A/k$, and thus $B | A/k$. The result follows. □

3 Main results

Let G be the Berlekamp matrix with respect to the polynomial $f(x)$ of degree m over F. The property $\mathcal{P}_1$ is defined as

$$\mathcal{P}_1 : \ G^m = I_m \quad \text{if and only if} \quad f(x) \text{ is irreducible}$$

where I_m is the identity matrix of order m.

Theorem 3.1. *Let $f(x)$ be a polynomial over F of degree $m \geq 2$. Then $f(x)$ has the property $\mathcal{P}_1$ if and only if*

$$m \text{ is an odd prime or } 9.$$

Proof. Note that $G^m = I_m$ if and only if $f(x)$ divides $x^{2^m} - x$. Then $\mathcal{P}_1$ is also realized as

$$\mathcal{P}_1' : \ f(x) \text{ divides } x^{2^m} - x \text{ if and only if } f(x) \text{ is irreducible.}$$

Moreover, since

$$x^{2^m} - x = \prod_{d|m} (\text{Irreducible polynomials of degree } d),$$

the property $\mathcal{P}_1'$ is equivalent to

$$\mathcal{P}_1'' : \ \text{If } f(x) \text{ divides } x^{2^m} - x \text{ then } f(x) \text{ is irreducible.}$$

To prove the sufficiency, suppose that $f(x)$ has the property $\mathcal{P}_1''$. Note that the number of irreducible polynomials $N(\ell)$ over F of degree ℓ is

$$N(\ell) \begin{cases} = 2 & \text{if } \ell = 1 \\ = \ell - 1 & \text{if } \ell = 2, 3, 4 \\ \geq \ell & \text{if } \ell > 4 \end{cases} .$$

Hence the degree m of $f(x)$ can not be written as $m = n\ell$ for some positive integers $1 < n \leq \ell$ and $\ell \geq 5$, or otherwise $N(\ell) \geq n$ and a product of n irreducible polynomials of degree ℓ does not have $\mathcal{P}_1''$. For the some reason, m can not be written as $m = n\ell$ for some positive integers $1 < n < \ell$, either. Furthermore, $m \neq 2$ and $m \neq 4$ since both of $x(x+1)$ and $x(x+1)(x^2+x+1)$ do not have $\mathcal{P}_1''$. To conclude the above argument, m is an odd prime or $m = 3 \cdot 3 = 9$.

For the necessity, assume m is an odd prime or 9. If m is an odd prime, then

$$x^{2^m} - x = x(x+1) \prod (\text{Irreducible polynomials of degree } m),$$

and hence $f(x)$ is an irreducible polynomial of degree m whenever $f(x)$ divides $x^{2^m} - x$. Besides, if $m = 9$ then

$$x^{2^9} - x = x(x+1)(x^3+x+1)(x^3+x^2+1) \prod (\text{Irreducible polynomials of degree } 9),$$

and thus $f(x)$ is an irreducible polynomial of degree 9 provided that $f(x)$ divides $x^{2^9} - x$. It says that $f(x)$ has $\mathcal{P}_1''$, and the proof is completed. $\square$

Let G be the Berlekamp matrix with respect to the polynomial $f(x)$ of degree m over F. The property $\mathcal{P}_2$ is defined as

$$\mathcal{P}_2 : \ o(G) = m \quad \text{if and only if} \quad f(x) \text{ is irreducible}$$

where $o(G)$ is the order of G.

Theorem 3.2. *Let $f(x)$ be a polynomial over F of degree $m \geq 2$. If $f(x)$ has the property $\mathcal{P}_2$ then m can be written as*

$$m = p^i \qquad or \qquad m = p^i q$$

for primes $p < q$ and positive integer i.

Proof. Suppose to the contrary that m can be written as a product of three pairwise coprime factors that more than 1, or $m = p^i q^j$ for primes $p < q$ and positive integers i, j with $j \geq 2$.

Assume that m can be written as $m = p_1 p_2 p_3$, where positive integers $1 < p_1 < p_2 < p_3$ are pairwise coprime. Since

$$p_1 p_2 p_3 = 1 \cdot p_1 p_3 + (p_2 - p_1) \cdot p_2 + (p_1 - 1) \cdot p_2 p_3,$$

a product of 1, $p_2 - p_1$, and $p_1 - 1$ irreducible polynomials respectively of degrees $p_1 p_3$, p_2, and $p_2 p_3$ does not have the property $\mathcal{P}_2$, which is a contradiction. (It is quick to check that the number $N_2(\ell)$ of irreducible polynomials over F of degree ℓ satisfies $N_2(\ell) \geq \ell - 1$ for each positive integer ℓ, and $\text{lcm}(p_1 p_3, p_2, p_2 p_3) = p_1 p_2 p_3 = m$.)

Next, suppose that m can be written as $m = p^i q^j$ for primes $p < q$ and positive integers i, j with $j \geq 2$. Since

$$p^i q^j = (p - 1) \cdot p^{i-1} q^j + 1 \cdot p^i q^{j-1} + (q - p) \cdot p^{i-1} q^{j-1},$$

a product of $p - 1$, 1, and $q - p$ irreducible polynomials respectively of degrees $p^{i-1} q^j$, $p^i q^{j-1}$, and $p^{i-1} q^{j-1}$ does not have the property $\mathcal{P}_2$, which is a contradiction. (It is immediate to check that $\text{lcm}(p^{i-1} q^j, p^i q^{j-1}, p^{i-1} q^{j-1}) = p^i q^j = m$.) The result follows. □

The case will be trivial if the degree m is a prime power. It may be quick to show that $f(x) \in F[x]$ has $\mathcal{P}_2$ if its degree $m = p^i$ for some prime p and positive integer i. Now, focus on the case $m = p^i q$ for primes $p < q$ and positive integer i.

Note that the order $o(f(x))$ of $f(x)$ is the least common multiple (l.c.m.) of the degrees of factors in $f(x)$. Then a quick observation is given below.

Lemma 3.3. *Assume that $f(x) \in F[x]$ is of degree $m = p^i q$ for primes $p < q$ and positive integer i. If the order $o(f(x)) = m$ then there exists a factor of $f(x)$ that of degree p^i or $p^i q$.*

□

Corollary 3.4. *Let $f(x) \in F[x]$ of degree $m = p^i q$ for primes $p < q$ and positive integer i. Then the following properties follow.*

(i) If $f(x)$ has $\mathcal{P}_2$ then a polynomial $\tilde{f}(x) \in F[x]$ of degree $\tilde{m} = p^{\tilde{i}} q$ with positive integer $\tilde{i} \leq i$ also has $\mathcal{P}_2$.

(ii) If $q > 2^{p^i}$ then $f(x)$ has $\mathcal{P}_2$.

(iii) If $p^i = 2$ then $q > 4$ if and only if $f(x)$ has $\mathcal{P}_2$.

(iv) If $q > p^i > 2$ and $f(x)$ has $\mathcal{P}_2$, then $(p^i - 2)q > 2^{p^i} - 2^{p^{i-1}} + 1$. If $q < p^i$ and $f(x)$ has $\mathcal{P}_2$, then $(q-2)p^i > 2^q$.

Proof. (i) is direct from Lemma 2.2.

To prove (ii), suppose $q > 2^{p^i}$ and the order $o(f(x))$ of $f(x)$ equals $m = p^i q$. Since $f(x)$ is of degree m and divides

$$x^{2^m} - x = (x^{2^{p^i}} - x) \prod_{j=1}^{i} (\text{Irreducible polynomials of degree } qj),$$

the degree of each factor of $f(x)$ is a multiple of q. Hence by Lemma 3.3 there is a factor in $f(x)$ that of degree $p^i q$, which means that $f(x)$ is composed of exactly one factor polynomial and is irreducible.

The sufficiency of (iii) is straightforward from (ii). Then the necessity of (iii) is only to examine the case $q = 3$, *i.e.*, $m = 2 \cdot 3 = 6$. One can see that the reducible polynomial $f(x) = x(x^2 + x + 1)(x^3 + x + 1)$ is of order $o(f(x)) = 6$, and thus does not have $\mathcal{P}_2$. The proof of (iii) is completed.

To prove the first part of (iv), suppose to the contrary that $q > p^i > 2$ and $f(x)$ has $\mathcal{P}_2$, but $(p^i - 2)q \leq 2^{p^i} - 2^{p^{i-1}} + 1$. Since $p < q$ are two distinct primes, one has $\gcd(p^i, q) = 1$, and thus

$$\{uq \pmod{p^i} \mid u = 1, 2, \ldots, p^i - 1\} = \{1, 2, \ldots, p^i - 1\}.$$

By Pigeon Hole Principle, there exists positive integer $\hat{u}$ with $1 \leq u \leq p^i - 2$ such that $\hat{u}q \equiv 1$ or $2 \pmod{p^i}$. If $\hat{u}q \equiv 1 \pmod{p^i}$, say, $\hat{u}q = \ell p^i + 1$, then the product of x and ℓ irreducible polynomials of degree p^i and $p^i - \hat{u}$ irreducible polynomials of degree q does not has $\mathcal{P}_2$, which is a contradiction. (Note that the fact $N(p^i) = (2^{p^i} - 2^{p^{i-1}})/p^i$ and $(p^i - 2)q \leq 2^{p^i} - 2^{p^{i-1}} + 1$ implies ℓ exists.) On the other hand, if $\hat{u}q \equiv 2 \pmod{p^i}$, say, $\hat{u}q = \ell p^i + 2$, then the product of $x(x+1)$ and ℓ irreducible polynomials of degree p^i and $p^i - \hat{u}$ irreducible polynomials of degree q does not has $\mathcal{P}_2$, which makes a contradiction. Similarly, to show the second part of (iv), suppose to the contrary that $q < p^i$ and $f(x)$ has $\mathcal{P}_2$, but $(q-2)p^i \leq 2^q$. An analogue version of the above contradiction will occur by exchanging the positions of p^i and q. The result follows. □

References

[1] E. R. Berlekamp, Factoring polynomials over finite fields, *Bell Syst. Tech. J.* 46 (1967) 1853-1859.

[2] J. von zur Gathen and D. Panario, Factoring polynomials over finite fields: A survey, *J. Symbolic Computation* 31 (2001) 3-17.

[3] K. Petr, Über die Reduzibilität eines Polynoms mit ganzzahligen Koeffizienten nach einem Primzahlmodul, *Časopis pro Pěstování matematiky a fysiky* 66 (1937) 85-94.

[4] R. G. Swan, Factorization of polynomials over Finite Fields, *Pacific J. Math.* 12 (1962) 1099-1106.